

RIVERINA



SALE & SHOW PELLETS / MEAL

PRODUCT DESCRIPTION

RIVERINA SALE & SHOW PELLETS / MEAL is a higher energy feed for stud cattle without urea and MUST be fed in conjunction with quality roughage.

TYPICAL COMPOSITION

- Barley, wheat, maize, sorghum, bran and pollard, cottonseed meal, full fat soybean meal, molasses, vegetable oil, limestone, sodium bentonite, dicalcium phosphate, buffer, potassium chloride, ammonium sulphate, salt, Rumensin®, flavouring, organic zinc, mould inhibitor, Riverina ruminant vitamin & mineral premix.
- Base raw materials may gradually change seasonally.

DIRECTIONS FOR USE

- **THIS IS NOT A COMPLETE FEED.** It has been designed to be fed with good quality hay and/or pasture feeding. As this feed contains higher starch levels more care should be taken with the introduction program.
- RIVERINA SALE & SHOW PELLETS / MEAL can be fed at 0.5-1.5% bodyweight/head/day using:
 - **Troughs** – split amount to be fed between 2 feeds. It is critical to have enough trough space for all animals to eat at the same time.
 - **Self-Feeders** – The slides need to be adjusted down to a finger width to control intakes. Gradually increase slide width to allow more access to the feed over a 2 month period. Feeders should be placed at the opposite end of the paddock to water troughs to promote grazing. Feed intakes should be monitored especially when commencing to feed.
 - **Partial Mixed Ration** – Can be mixed with silage/hay and other ingredients to form a complete or partial mixed ration to be fed in a semi feedlot situation.

Typical Program:

- Start background feeding 6 months prior to sale. Commence feeding at 0.5% of body weight. This can be gradually increased over a couple of months to 1.0% of body weight.
- 2 months prior to sale the cattle can be put into yards to finish them. At this time feeding levels can be increased to 1.5% of body weight and fed in conjunction with good quality grassy hay supplied *ad lib* in hay racks.
- Ensure that there is continuous access to fresh, cool, clean water.
- When feeding cattle, high grain feeds should be gradually introduced, to allow the rumen to adapt and to avoid acidosis/ grain poisoning. For stud cattle it is recommended that this be done over at least 2 months.

NOTE: There are many factors involved in the preparation of stud cattle for sale, so contact your Riverina Representative for more complete advice on how to feed your stud cattle.

NUTRITIONAL ANALYSIS

PROTEIN	%	MIN:	14.50
UREA	%		Nil
CALCIUM	%	:	0.70 - 1.00
PHOSPHORUS	%	:	0.30 - 0.70
C FIBRE	%	MIN:	6.00
ENERGY ME	MJ/kg	MIN:	11.30

NUTRITIONAL ANALYSIS (continued)

SALT	%	:	0.50
VITAMIN A	iu/kg	:	6,700.00
VITAMIN D3	iu/kg	:	800.00
VITAMIN E	mg/kg	:	25.00
IRON	mg/kg	:	50.00
IODINE	mg/kg	:	0.50
COBALT	mg/kg	:	0.50
COPPER	mg/kg	:	12.00
MANGANESE	mg/kg	:	40.00
SELENIUM	mg/kg	:	0.10
ZINC	mg/kg	:	72.00

FEATURES

- Organic zinc to aid hoof condition.
- Flavour is added to improve acceptance by cattle.
- RIVERINA SALE & SHOW PELLETS / MEAL includes rumen buffers and modifiers to enhance performance.
- Oil in the added full fat soybean meal will help improve coat condition.
- High by pass protein levels with NO Urea added.

QUALITY PRODUCT

- Manufactured to strict quality control standards for incoming raw materials and the finished product.
- Formulated in accordance with current research into ruminant nutrition and mixed using a variety of quality raw materials.

MEDICATION

This feed contains Rumensin® which supplies Monensin sodium at 30 mg/kg to improve feed efficiency and weight gain (pasture cattle).

WITHOLDING PERIOD

Nil.

WARNING

**Do not feed to dogs, horses or other equines as it may prove fatal.
Do not feed to sheep.
Feed only to cattle.**

STORAGE

[Cool, shaded, dry conditions, away from vermin.](#)

EXPIRY

[Six months from the date of manufacture.](#)

PACK SIZE

20kg woven polypropylene bags.
Bulk: Please contact your Riverina Sales Representative.



Rumensin® is a registered trademark of Eli Lilly and Company.

This feed is scientifically formulated for the intended species of animal and should not be fed to any other animal. It may contain medication and ingredients that may prove harmful if fed to other species.
Ver: 7.0

Issue Date: October 2015

